

兔抗 BCAM 多克隆抗体

中文名称：兔抗 BCAM 多克隆抗体

英文名称：Anti-BCAM rabbit polyclonal antibody

别名：AU; LU; CD239; MSK19

相关类别：一抗

抗原：BCAM

储存：冷冻（-20℃）

宿主：Rabbit

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:

This gene encodes Lutheran blood group glycoprotein, a member of the immunoglobulin superfamily and a receptor for the extracellular matrix protein, laminin. The protein contains five extracellular immunoglobulin domains, a single transmembrane domain, and a short C-terminal cytoplasmic tail. This protein may play a role in epithelial cell cancer and in vaso-occlusion of red blood cells in sickle cell disease. Polymorphisms in this gene define some of the antigens in the Lutheran system and also the Auberger system. Inactivating variants of

| | |
|------------------------------------|--|
| | this gene result in the recessive Lutheran null phenotype, Lu(a-b-), of the Lutheran blood group. Two transcript variants encoding different isoforms have been found for this gene. |
| Applications: | ELISA, WB |
| Name of antibody: | BCAM |
| Immunogen: | Fusion protein of human BCAM |
| Full name: | basal cell adhesion molecule (Lutheran blood group) |
| Synonyms : | AU; LU; CD239; MSK19 |
| SwissProt: | P50895 |
| ELISA Recommended dilution: | 1000-2000 |
| WB Predicted band size: | 67 kDa |
| WB Positive control: | 293T and 231 cells |
| WB Recommended dilution: | 200-1000 |

