

## 兔抗 CD160 多克隆抗体

中文名称: 兔抗 CD160 多克隆抗体

英文名称: Anti-CD160 rabbit polyclonal antibody

别 名: NK1; BY55; NK28

相关类别: 一抗

抗原: CD160

储 存: 冷冻(-20℃)

宿 主: Rabbit

反应种属: Human

标记物: Unconjugate

克隆类型: rabbit polyclonal

## 技术规格

**Background:** 

CD160 is an 27 kDa glycoprotein which was initially ide ntified with the monoclonal antibody BY55. Its expressi on is tightly associated with peripheral blood NK cells and CD8 T lymphocytes with cytolytic effector activity. The cDNA sequence of CD160 predicts a cysteine-rich, glycosylphosphatidylinositol-anchored protein of 181 am ino acids with a single Ig-like domain weakly homologo us to KIR2DL4 molecule. CD160 is expressed at the cell surface as a tightly disulfide-linked multimer. RNA blot analysis revealed CD160 mRNAs of 1.5 and 1.6 kb who se expression was highly restricted to circulating NK an



	d T cells, spleen and small intestine.
Applications:	ELISA, WB, IHC
Name of antibody:	CD160
Immunogen:	Fusion protein of human CD160
Full name:	CD160 molecule
Synonyms:	NK1; BY55; NK28
SwissProt:	O95971
ELISA Recommended dilution:	2000-5000
IHC positive control:	Human thyroid cancer and human gastric cancer
IHC Recommend dilution:	25-100
WB Predicted band size:	20 kDa
WB Positive control:	Hela cells
WB Recommended dilution:	500-2000





