

兔抗 CDK11B 多克隆抗体

中文名称: 兔抗 CDK11B 多克隆抗体

英文名称: Anti-CDK11B rabbit polyclonal antibody

别 名: p58; PK58; CDK11; CLK-1; CDC2L1; PITSLREA; p58CLK-1; CDK11-p46; CDK11-p58; p58CDC2L1; CDK11-p110

相关类别: 一抗

储 存: 冷冻(-20℃) 避光

抗 原: CDK11B

宿 主: Rabbit

反应种属: Human Mouse

标记物: Unconjugate

克隆类型: Unconjugate

技术规格

Applications:	WB
	may play a role in cell apoptosis.
	oded by this gene can be cleaved by caspases and
	or altered in neuroblastoma. The protein kinase enc
	oring gene. These two genes are frequently deleted
Background:	e shares very high sequence identity with a neighb
	le control. Due to a segmental duplication, this gen
	ily are known to be essential for eukaryotic cell cyc
	e protein kinase family. Members of this kinase fam
	This gene encodes a member of the serine/threonin



Name of antibody:	CDK11B
Immunogen:	Synthesized peptide derived from N-terminal of hu man CD2L1.
Full name:	cyclin-dependent kinase 11B
Synonyms :	p58; PK58; CDK11; CLK-1; CDC2L1; PITSLREA; p58CL K-1; CDK11-p46; CDK11-p58; p58CDC2L1; CDK11-p1 10
SwissProt:	P21127
WB Predicted band size:	93 kDa
WB Positive control:	HepG2 cells and Huvec cells lysates
WB Recommended dilution:	500-3000

