

兔抗 NR4A3 多克隆抗体

- 中文名称: 兔抗 NR4A3 多克隆抗体
- 英文名称: Anti-NR4A3 rabbit polyclonal antibody
- 别 名: CHN; TEC; CSMF; NOR1; MINOR
- 储 存: 冷冻 (-20℃) 避光
- 抗 原: NR4A3
- 宿 主: Rabbit
- 相关类别: 一抗
- 反应种属: Human
- 标记物: Unconjugate
- 克隆类型: Unconjugate
- 技术规格

	This gene encodes a member of the steroid-thyroid hormo
	ne-retinoid receptor superfamily. The encoded protein may
	act as a transcriptional activator. The protein can efficiently
	bind the NGFI-B Response Element (NBRE). Three different
	versions of extraskeletal myxoid chondrosarcomas (EMCs) a
	re the result of reciprocal translocations between this gene
Background:	and other genes. The translocation breakpoints are associat
	ed with Nuclear Receptor Subfamily 4, Group A, Member 3
	(on chromosome 9) and either Ewing Sarcome Breakpoint
	Region 1 (on chromosome 22), RNA Polymerase II, TATA B
	ox-Binding Protein-Associated Factor, 68-KD (on chromoso
	me 17), or Transcription factor 12 (on chromosome 15). M
	ultiple transcript variants encoding different isoforms have



	been found for this gene.
Applications:	WB
Name of antibody:	NR4A3
Immunogen:	Synthesized peptide derived from internal of human NR4A 3.
Full name:	nuclear receptor subfamily 4, group A, member 3
Synonyms :	CHN; TEC; CSMF; NOR1; MINOR
SwissProt:	Q92570
WB Predicted band size:	68 kDa
WB Positive control:	COLO cells lysate
WB Recommended dilution:	500-3000

