

小鼠抗 UBE2J1 单克隆抗体

中文名称: 小鼠抗 UBE2J1 单克隆抗体

英文名称: Anti-UBE2J1 mouse monoclonal antibody

别名: CGI-76; HSPC153; HSPC205; HSU93243; NCUBE-1; NCUBE1; UBC6; UBC6E; Ubc6p

储存: 冷冻 (-20℃) 避光

抗原: UBE2J1

宿主: Mouse

反应种属: Human , Dog , Rat , Monkey , Mouse

相关类别: 一抗

标记物: Unconjugate

克隆类型: mouse monoclonal

技术规格

Background:	The modification of proteins with ubiquitin is an important cellular mechanism for targeting abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes, or E1s, ubiquitin-conjugating enzymes, or E2s, and ubiquitin-protein ligases, or E3s. This gene encodes a member of the E2 ubiquitin-conjugating enzyme family. This enzyme is located in the membrane of the endoplasmic reticulum (ER) and may contribute to quality control ER-associated degradation by the ubiquitin-proteasome system. [provided by RefSeq]. COMPLETENESS: complete on the 3' end.
Applications:	WB, IHC, FC
Name of antibody:	UBE2J1

Immunogen:	Fusion protein of human UBE2J1
Full name:	ubiquitin-conjugating enzyme E2, J1 (UBE2J1)
Synonyms:	CGI-76; HSPC153; HSPC205; HSU93243; NCUBE-1; NCUBE1; UBC6; UBC6E; Ubc6p
SwissProt:	Q9Y385
IHC positive control:	carcinoma of human lung tissue and carcinoma of human prostate tissue; human liver tissue and carcinoma of human liver tissue
IHC Recommend dilution:	30-150
WB Predicted band size:	35 kDa
WB Positive control:	HepG2, HeLa, SVT2, A549, COS7, Jurkat, MDCK, PC12, MCF-7 cell lysates
WB Recommended dilution:	500-2000