

SEPTIN2 抗原（重组蛋白）

中文名称： SEPTIN2 抗原（重组蛋白）

英文名称： SEPTIN2 Antigen (Recombinant Protein)

别名： septin 2; DIFF6; NEDD5; SEPT2; NEDD-5; Pnutl3; hNedd5

储存： 冷冻（-20℃）

相关类别： 抗原

概述

Fusion protein corresponding to a region derived from 1-371 amino acids of human SEPTIN2

技术规格

Full name:	septin 2
Synonyms:	DIFF6; NEDD5; SEPT2; NEDD-5; Pnutl3; hNedd5
Swissprot:	Q15019
Gene Accession:	BC033559
Purity:	>85%, as determined by Coomassie blue stained SDS-PAGE
Expression system:	Escherichia coli
Tags:	His tag C-Terminus, GST tag N-Terminus
Background:	Filament-forming cytoskeletal GTPase. Forms a filamentous structure with SEPTIN12, SEPTIN6, SEPTIN2 and probably SEPTIN4 at the sperm annulus which is required for the structural integrity and motility of the sperm tail during post meiotic differentiation (PubMed:25588830). Required for normal organization of the actin cytoskeleton. Plays a role i

in the biogenesis of polarized columnar-shaped epithelium by maintaining polyglutamylated microtubules, thus facilitating efficient vesicle transport, and by impeding MAP4 binding to tubulin. Required for the progression through mitosis. Forms a scaffold at the midplane of the mitotic spindle required to maintain CENPE localization at kinetochores and consequently chromosome congression. During anaphase, may be required for chromosome segregation and spindle elongation. Plays a role in ciliogenesis and collective cell movements. In cilia, required for the integrity of the diffusion barrier at the base of the primary cilium that prevents diffusion of transmembrane proteins between the cilia and plasma membranes: probably acts by regulating the assembly of the tectonic-like complex (also named B9 complex) by localizing TMEM231 protein. May play a role in the internalization of 2 intracellular microbial pathogens, *Listeria monocytogenes* and *Shigella flexneri*.