

兔抗 CDK2(Phospho-Thr160)多克隆抗体

中文名称：兔抗 CDK2(Phospho-Thr160)多克隆抗体

英文名称：Anti-CDK2(Phospho-Thr160) polyclonal antibody

别名：cyclin dependent kinase 2; CDKN2; p33(CDK2)

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：CDK2(Phospho-Thr160)

反应种属：Human Mouse Rat

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:	This gene encodes a member of a family of serine/threonine protein kinases that participate in cell cycle regulation. The encoded protein is the catalytic subunit of the cyclin-dependent protein kinase complex, which regulates progression through the cell cycle. Activity of this protein is especially critical during the G1 to S phase transition. This protein associates with and regulated by other subunits of the complex including cyclin A or E, CDK inhibitor p21Cip1 (CDKN1A), and p27Kip1 (CDKN1B)
Applications:	WB, IHC
Name of antibody:	CDK2(Phospho-Thr160)

Immunogen:	Peptide sequence around phosphorylation site of threonine 160 (T-Y-T(p)-H-E) derived from Human CDK2
Full name:	cyclin-dependent kinase 2
Synonyms:	CDKN2; p33(CDK2)
SwissProt:	P24941
IHC positive control:	Human breast carcinoma
IHC Recommend dilution:	50-100
WB Predicted band size:	34 kDa
WB Positive control:	Hela cells treated with UV
WB Recommended dilution:	500-1000



